HER2陰性乳癌
(Luminal A, triple negative)
に対する
術後化学療法レジメン

相良病院

<背景>

- 医師の化学療法レジメンに関する見解に、若干の相違がある
- 朝のカンファで毎回、ディスカッションが白熱してしまい、外来診療の開始に支障を来している

<目的>

- 現在までのエビデンスをもとに、化学療法レジメンに関する、基本的な院内コンセンサス作り

症例1 60歳 女性

- Breast (Rt):invasive ductal carcinoma, a3
- f, pT2, ly+, v-, pn-, wide excision
- Lymph node: no metasis, 2/12, リンパ節転移巣5mm sentinel with axillary dissection
- ER: 90%以上, PgR: 50-90%, Hercep test: スコア1+
- 核異型度: Grade 2
- 組織学的腫瘍の広がり 50x35mm、浸潤径 25x15mm

Luminal A, node positive, リスク因子 3個

症例2 39歳 女性

- Breast (Rt):invasive ductal carcinoma, a2
- g, pT1c, ly-, v-, pn-, wide excision
- Lymph node: no metasis, 0/3, sentinel sampling
- ER: 1-10%, PgR: <1%, Hercep test: スコア1+
- 核異型度 : Grade 3
- 組織学的腫瘍の広がり 26x10mm、浸潤径 25x10mm

Triple negative, node negative, リスク因子 3個

- AC4 ?
- FEC 6 ?
- AC → Taxane ?
- FEC → Taxane?

AC or FEC ?

Anthracycline Strength

EBCTCG criteria

Type of Anthracycline	Dose Intensity	Cumulative Dose	Schedule of Administration
Doxorubicin	$\geq 25\text{mg/m}^2/\text{week}$	$\geq 240\text{ mg/m}^2$	Day 1 + Day 8 Q3-4 weeks
Epirubicin	$\geq 30\text{mg/m}^2/\text{Week}$	$\geq 360\text{ mg/m}^2$	Day 1 + Day 8 Q3-4 weeks

Standard Strength = ≥ 2 criteria

Lower Strength = < 2 criteria

Anthracycline Strength

Standard Strength

FEC100 x 4 - 6

FEC90 x 4 - 6

CEF x 4 - 6

CAF x 4 - 6

A75 or E100 x 4 → CMF

Lower Strength

FEC75 x 6

FEC60 x 6

FEC50 x 6

FAC 50 x 6

AC x 4

EC x 4

Relapse free survival MA5

Fig 1. Ten-year relapse-free survival of cyclophosphamide, methotrexate, and fluorouracil (CMF) versus cyclophosphamide, epirubicin, and fluorouracil (CEF).

CEF 6cycle

C: 75 mg/m² days 1 ~ 14,

E: 60 mg/m² **days 1 and 8**

F: 500 mg/m² days 1 and 8

CMF 6cycle

C: 100 mg/m² days 1 ~ 14

M: 40 mg/m² days 1 and 8

F: 600 mg/m² days 1 and 8

NSABP B-30

Overall Survival (Intention-To-Treat)

CMF = AC 4 = AT 4 = TAC 4

どのような症例に
Taxaneの追加を行うか？

リンパ節転移個数とタキサン追加効果

ホルモン受容体とタキサン追加効果

Disease Free Survival – ER/HER2 subgroups Compared with other trials

Hazard ratios by
ER & HER2 status

Sequential docetaxel as adjuvant chemotherapy for early breast cancer (TACT): an open-label, phase III, randomised controlled trial

Paul Ellis, Peter Barrett-Lee, Lindsay Johnson, David Cameron, Andrew Wardley, Susan O'Reilly, Mark Verrill, Ian Smith, John Yarnold, Robert Coleman, Helena Earl, Peter Canney, Chris Twelves, Christopher Poole, David Bloomfield, Penelope Hopwood, Stephen Johnston, Mitchell Dowsett, John M S Bartlett, Ian Ellis, Clare Peckitt, Emma Hall, Judith M Bliss, for the TACT Trial Management Group and the TACT Trialists*

Summary

Background Incorporation of a taxane as adjuvant treatment for early breast cancer offers potential for further improvement of anthracycline-based treatment. The UK TACT study (CRUK01/001) investigated whether sequential docetaxel after anthracycline chemotherapy would improve patient outcome compared with standard chemotherapy of similar duration.

Methods In this multicentre, open-label, phase III, randomised controlled trial, 4162 women (aged >18 years) with node-positive or high-risk node-negative operable early breast cancer were randomly assigned by computer-generated permuted block randomisation to receive FEC (fluorouracil 600 mg/m², epirubicin 60 mg/m², cyclophosphamide 600 mg/m² at 3-weekly intervals) for four cycles followed by docetaxel (100 mg/m² at 3-weekly intervals) for four cycles (n=2073) or control (n=2089). For the control regimen, centres chose either FEC for eight cycles (n=1265) or epirubicin (100 mg/m² at 3-weekly intervals) for four cycles followed by CMF (cyclophosphamide 600 mg/m², methotrexate 40 mg/m², and fluorouracil 600 mg/m² at 4-weekly intervals) for four cycles (n=824). The primary endpoint was disease-free survival. Analysis was by intention to treat (ITT). This study is registered as an International Standard Randomised Controlled Trial, number ISRCTN79718493.

Findings All randomised patients were included in the ITT population. With a median follow-up of 62 months, disease-free survival events were seen in 517 of 2073 patients in the experimental group compared with 539 of 2089 controls (hazard ratio [HR] 0.95, 95% CI 0.85–1.08; p=0.44). 75.6% (95% CI 73.7–77.5) of patients in the experimental group and 74.3% (72.3–76.2) of controls were alive and disease-free at 5 years. The proportion of patients who reported any acute grade 3 or 4 adverse event was significantly greater in the experimental group than in the control group (p<0.0001); the most frequent events were neutropenia (937 events vs 797 events), leucopenia (507 vs 362), and lethargy (456 vs 272).

Interpretation This study did not show any overall gain from the addition of docetaxel to standard anthracycline chemotherapy. Exploration of predictive biomarker-defined subgroups might have the potential to better target the use of taxane-based therapy.

Lancet 2009; 373: 1681–92

See [Comment](#) page 1662

*Members listed at end of paper

Guy's and St Thomas' NHS Trust, London, UK (P Ellis MD); Velindre NHS Trust, Cardiff, UK (Prof P Barrett-Lee MD); The Institute of Cancer Research, Sutton, UK (L Johnson BSc, C Peckitt MSc, E Hall PhD, Prof J M Bliss MSc); University of Leeds, Leeds, UK (Prof D Cameron MD); Christie Hospital NHS Trust, Manchester, UK (A Wardley MD, Prof P Hopwood MD); Clatterbridge Centre for Oncology, Bebington, UK (S O'Reilly MD); Newcastle upon Tyne Hospitals NHS Trust, Newcastle, UK (M Verrill FRCP); Royal Marsden NHS Trust, London and Sutton, UK (Prof I Smith MD, Prof J Yarnold FRCP, S Johnston FRCP, Prof M Dowsett PhD); Weston Park Hospital, Sheffield, UK (Prof R Coleman MD); Addenbrooke's Hospital, Cambridge, UK (H Earl FRCP); Beatson Centre for Oncology,

Number of events/ number at risk		0	1	2	3	4	5
Control	0/2089	81/2001	154/1838	121/1709	91/1537	68(+24)/888	
FEC-D	0/2073	62/2006	145/1859	141/1708	88/1541	52(+29)/905

Number of events/ number at risk		0	1	2	3	4	5
Control	0/2089	25/2057	77/1971	98/1864	81/1695	57(+40)/1000	
FEC-D	0/2073	22/2046	71/1972	101/1861	67/1708	82(+31)/993	

FEC4-D4 (n=2073)

F:600 mg/m²
E:60 mg/m²,
C: 600 mg/m²
at 3-weekly intervals
D: 100 mg/m²
at 3-weekly intervals

Control

FEC8 (n=1265)

F:600 mg/m²
E:60 mg/m²,
C: 600 mg/m²
at 3-weekly intervals

FEC4-CMF4 (n=824)

E: 100 mg/m²
at 3-weekly intervals

エンドキサンによる
Ovarian suppression効果？
E:100の効果？

Figure 5: Hazard ratios for disease-free survival by combined ER and HER2 status in (A) all patients and (B) node-positive patients

ER=oestrogen receptor. FEC-D=fluorouracil, epirubicin, and cyclophosphamide followed by docetaxel.

HER2=human epidermal growth factor receptor 2. HR=hazard ratio.

Randomized Phase 3 Trial of Fluorouracil, Epirubicin, and Cyclophosphamide Alone or Followed by Paclitaxel for Early Breast Cancer

Miguel Martín, Álvaro Rodríguez-Lescure, Amparo Ruiz, Emilio Alba, Lourdes Calvo, Manuel Ruiz-Borrego, Blanca Munárriz, César A. Rodríguez, Carmen Crespo, Enrique de Alava, José Antonio López García-Asenjo, María Dolores Guitián, Sergio Almenar, Jesús Fernando González-Palacios, Francisco Vera, José Palacios, Manuel Ramos, Jose Manuel Gracia Marco, Ana Lluch, Isabel Alvarez, Miguel Angel Seguí, José Ignacio Mayordomo, Antonio Antón, José Manuel Baena, Arrate Plazaola, Alfonso Modolell, Amadeu Pelegrí, Jose Ramón Mel, Enrique Aranda, Encarna Adrover, José Valero Álvarez, José Luis García Puche, Pedro Sánchez-Rovira, Sonia Gonzalez, José Manuel López-Vega

On behalf of the **GEICAM** 9906 Study Investigators

- Background** Taxanes are among the most active drugs for the treatment of metastatic breast cancer, and, as a consequence, they have also been studied in the adjuvant setting.
- Methods** After breast cancer surgery, women with lymph node–positive disease were randomly assigned to treatment with fluorouracil, epirubicin, and cyclophosphamide (FEC) or with FEC followed by weekly paclitaxel (FEC-P). The primary endpoint of study—5-year disease-free survival (DFS)—was assessed by Kaplan–Meier analysis. Secondary endpoints included overall survival and analysis of the prognostic and predictive value of clinical and molecular (hormone receptors by immunohistochemistry and HER2 by fluorescence in situ hybridization) markers. Associations and interactions were assessed with a multivariable Cox proportional hazards model for DFS for the following covariates: age, menopausal status, tumor size, lymph node status, type of chemotherapy, tumor size, positive lymph nodes, HER2 status, and hormone receptor status. All statistical tests were two-sided.
- Results** Among the 1246 eligible patients, estimated rates of DFS at 5 years were 78.5% in the FEC-P arm and 72.1% in the FEC arm (difference = 6.4%, 95% confidence interval [CI] = 1.6% to 11.2%; $P = .006$). FEC-P treatment was associated with a 23% reduction in the risk of relapse compared with FEC treatment (146 relapses in the 614 patients in the FEC-P arm vs 193 relapses in the 632 patients in the FEC arm, hazard ratio [HR] = 0.77, 95% CI = 0.62 to 0.95; $P = .022$) and a 22% reduction in the risk of death (73 and 95 deaths, respectively, HR = 0.78, 95% CI = 0.57 to 1.06; $P = .110$). Among the 928 patients for whom tumor samples were centrally analyzed, type of chemotherapy (FEC vs FEC-P) ($P = .017$), number of involved axillary lymph nodes ($P < .001$), tumor size ($P = .020$), hormone receptor status ($P = .004$), and HER2 status ($P = .006$) were all associated with DFS. We found no statistically significant interaction between HER2 status and paclitaxel treatment or between hormone receptor status and paclitaxel treatment.
- Conclusions** Among patients with operable breast cancer, FEC-P treatment statistically significantly reduced the risk of relapse compared with FEC as adjuvant therapy.

J Natl Cancer Inst 2008;100:805–814

[FEC-P arm \(n=632\)](#)

- FEC
at 3weekly, 4cycle

- Paclitaxe
P:100mg/m²
at weekly, 8cycle

[FEC arm \(n=614\)](#)

- F: 600mg/m²

- E: 90mg/m²

- C: 600mg/m²

- at 3weekly, 6 cycle

Ki 67とdocetaxelの追加効果との関係

PACS 01

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0	E C→E	E C→E	C→E E	C→E E	C
	n=1-3	E C→E	E C→E	C→E E	C→E E	C
高リスク	n=1-3	C→E	C→E	C→E	C→E	C
	n≥4	C→E	C→E	C→E	C→E	C

化学療法の適応は？

化学療法の
メリットが少ない

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

化学療法の適応は？

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0	E C→E	E C→E	C→E E	C→E E	C
	n=1-3	E C→E	E C→E	C→E E	C→E E	C
高リスク	n=1-3	C→E	C→E	C→E	C→E	C
	n≥4	C→E	C→E	C→E	C→E	C

oncotype DX[®]
Breast Cancer Assay

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0	E C→E	E C→E	C→E E	C→E E	C
	n=1-3	E C→E	E C→E	C→E E	C→E E	C
高リスク	n=1-3	C→E	C→E	C→E	C→E	C
	n≥4	C→E	C→E	C→E	C→E	C

化学療法の適応は？

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0	E C→E	E C→E	C→E E	C→E E	C
	n=1-3	E C→E	E C→E	C→E E	C→E E	C
高リスク	n=1-3	C→E	C→E	C→E	C→E	C
	n≥4	C→E	C→E	C→E	C→E	C

化学療法のレジメンは？

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0					
	n=1-3					
高リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
リンパ節転移個数を重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間 リスク	n=0					
	n=1-3					
高 リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0					
	n=1-3					
高リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0					
	n=1-3					
高リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0					
	n=1-3					
高リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間リスク	n=0					
	n=1-3					
高リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab

HER2		HER2陰性				
内分泌		反応性		不完全反応性		非反応性
閉経		pre	post	pre	post	Pre and post
低リスク		E	E	E	E	
中間 リスク	n=0					
	n=1-3					
高 リスク	n=1-3					
	n≥4					

化学療法のレジメンは？
Risk とresponseを重視すると・・・

化学療法レジメン

AC4 or TC4

AC4 → T

FEC → T

C: chemotherapy, E: endocrine therapy, Tr: Trastuzumab